BRITISH GEOLOGICAL SURVEY King Edward Point Observatory Monthly Magnetic Bulletin **December 2015** 15/12/KE









British Geological Survey NATURAL ENVIRONMENT RESEARCH COUNCIL

KING EDWARD POINT OBSERVATORY MAGNETIC DATA

1. Introduction

King Edward Point observatory was installed by the British Geological Survey (BGS), with the permission of the Government of South Georgia and the South Sandwich Islands (GSGSSI), in collaboration with British Antarctic Survey (BAS), and became operational in February 2011.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team Earth Hazards and Systems British Geological Survey Murchison House, West Mains Road Edinburgh EH9 3LA Scotland, UK

Tel:	+44 (0) 131 667 1000						
Fax:	+44 (0) 131 668 2683						
E-mail:	enquiries@bgs.ac.uk						
Internet: www.geomag.bgs.ac.uk							

2. Position

King Edward Point Observatory, one of the geomagnetic observatories maintained and operated by the British Geological Survey (BGS), is situated on a site adjacent to the Government of South Georgia and the South Sandwich Islands (GSGSSI) research station at King Edward Point, Cumberland East Bay, South Georgia. The observatory co-ordinates are:

 Geographic:
 54° 16'55.7"S
 323° 30'25.6"E

 Geomagnetic:
 46°09'47"S
 029° 50'42"E

 Height above mean sea level:
 7 m

The geographical coordinates are measured by a handheld GPS device, which uses WGS84 as the reference coordinate system. The height above MSL is determined from the best available contour maps. The geomagnetic co-ordinates are approximations, calculated using the 12th generation International Geomagnetic Reference Field (IGRF) at epoch 2015.5. On-line access to models (including IGRF), charts and navigational data are available at

http://www.geomag.bgs.ac.uk/data_service/models _compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became operational in February 2011. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at

www.geomag.bgs.ac.uk/data_service/home

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. One set of absolute measurements of the field are made manually once per month. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F. The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of

the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D, H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D, H and Z. The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using oneminute values of D, H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D, H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

4.5 Daily and Monthly Mean Values

Daily mean values of D, H, Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of Use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.

Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at King Edward Point magnetic observatory, South Georgia operated by the British Geological Survey in cooperation with the British Antarctic Survey and the Government of South Georgia and the South Sandwich Islands (GSGSSI).'


Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff


© NERC 2015. All rights reserved


KING EDWARD POINT OBSERVATORY


ABSOLUTE OBSERVATIONS


		Declination			Inclination		Total Field		Horizontal Intensity		Vertical Intensity		
Date	Day Number	Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	Observer
02-Dec-15	336	17:39	-7.1043	-7.0433	18:03	-55.7568	12.3	27882.7	15689.8	15998.0	-23049.4	-23198.9	LC
02-Dec-15	336	18:18	-7.1344	-7.0450	18:30	-55.7516	12.3	27882.5	15691.8	15997.7	-23047.8	-23199.1	LC
17-Dec-15	351	13:46	-7.1970	-7.0667	14:09	-55.7575	12.3	27883.2	15689.8	15997.8	-23050.0	-23199.1	LC
17-Dec-15	351	14:26	-7.1612	-7.0400	14:42	-55.7493	12.3	27883.6	15693.3	15997.8	-23048.1	-23199.1	LC
























































King Edward Point Observatory: Declination (degrees)




King Edward Point Observatory: Horizontal Intensity fhTŁ





King Edward Point Observatory: Vertical Intensity (nT)



Monthly Mean Values for King Edward Point Observatory 2015

Month	D	Н	Ι	Х	Y	Ζ	F
January February March April May June July August September October November	-7° 11.6′ -7° 11.9′ -7° 12.4′ -7° 12.4′ -7° 12.7′ -7° 12.9′ -7° 12.9′ -7° 12.9′ -7° 12.2′ -7° 12.0′ -7° 11.3′	15753 nT 15742 nT 15723 nT 15721 nT 15720 nT 15710 nT 15705 nT 15698 nT 15694 nT 15694 nT	-55° 40.9′ -55° 41.9′ -55° 44.0′ -55° 44.2′ -55° 44.8′ -55° 45.4′ -55° 45.8′ -55° 46.1′ -55° 46.1′ -55° 45.4′	15629 nT 15618 nT 15599 nT 15596 nT 15586 nT 15586 nT 15580 nT 15574 nT 15570 nT 15565 nT 15571 nT	-1972 nT -1973 nT -1972 nT -1973 nT -1972 nT -1972 nT -1973 nT -1972 nT -1968 nT -1967 nT -1964 nT	-23077 nT -23076 nT -23078 nT -23076 nT -23074 nT -23071 nT -23067 nT -23066 nT -23063 nT -23056 nT	27941 nT 27934 nT 27925 nT 27922 nT 27920 nT 27911 nT 27909 nT 27902 nT 27898 nT 27893 nT 27891 nT
December	-7° 11.5′	15692 nT	-55° 45.6′	15569 nT	-1965 nT	-23055 nT	27889 nT

Note

i. The values shown here are provisional.