

**BRITISH GEOLOGICAL SURVEY**

**Sable Island  
Observatory  
Monthly  
Magnetic  
Bulletin**

**January 2005**

**05/01/SB**



**British  
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

# SABLE ISLAND OBSERVATORY MAGNETIC DATA

## 1.1 Introduction

Sable Island is the third overseas geomagnetic observatory to be established by BGS. The installation, funded by a joint venture between BGS, Sperry Drilling Services and Sable Offshore Energy, was completed in May 1999 and the observatory became operational from 8<sup>th</sup> May 1999.

Magnetic observatory data is presented as a series of plots of one-minute, hourly and daily values, followed by a tabulation of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

National Geomagnetic Service  
British Geological Survey  
Murchison House, West Mains Road  
Edinburgh EH9 3LA  
Scotland, UK

Tel: +44 (0) 131 667 1000  
Fax: +44 (0) 131 668 4368  
E-mail: o.baillie@bgs.ac.uk  
Internet: [www.geomag.bgs.ac.uk](http://www.geomag.bgs.ac.uk)

## 1.2 Position

The Island is a sandbank formed by the meeting of currents from the St. Lawrence Delta and the Gulf Stream and is located approximately 290km southeast of Halifax, Nova Scotia.

The observatory co-ordinates are:-

Geographic:  $43^{\circ} 55.9'N$   $299^{\circ}0.4'E$   
Geomagnetic:  $52^{\circ} 07.6'N$   $11^{\circ} 54.4'E$   
Height above mean sea level:  $5m$  (approx)

The geomagnetic co-ordinates are calculated using the 10<sup>th</sup> generation International Geomagnetic Reference Field (IGRF) at epoch 2005.5

## 1.3 The Observatory Operation

### 1.3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed in April 2004 and became fully operational from 13<sup>th</sup> May 2004. The system operates under the control of data acquisition software running on QNX computers, which control the data logging and communications.

There are two sets of sensors used for making magnetic measurements. A triaxial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the

variations in the horizontal ( $H$ ) and vertical ( $Z$ ) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination ( $D$ ). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer making measurements of the absolute total field intensity ( $F$ ) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter whilst the total field intensity samples are filtered using a 7-point cosine filter.

## 1.4 Data Presentation

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

### 1.4.1 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days a page and show the variations in  $D$ ,  $H$  and  $Z$ . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

### 1.4.2 Magnetograms

The daily magnetograms are plotted using one-minute values of  $D$ ,  $H$  and  $Z$  from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.


### 1.4.3 Hourly Mean Value Plots


Hourly mean values of  $D$ ,  $H$  and  $Z$  for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions on the surface of the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence, and also illustrates seasonal and diurnal variations throughout the year.

### 1.4.4 Daily and Monthly Mean Values

Daily mean values of  $D$ ,  $H$ ,  $Z$  and  $F$  are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive.

Sable Island January 2005






Date: 01-01-2005


# Sable Island

Day number: 001



Date: 02-01-2005


Day number: 002



Date: 03-01-2005


# Sable Island

Day number: 003



Date: 04-01-2005


Day number: 004



Date: 05-01-2005


# Sable Island

Day number: 005



Date: 06-01-2005


Day number: 006



Date: 07-01-2005


# Sable Island

Day number: 007



Date: 08-01-2005

Day number: 008






Date: 09-01-2005


# Sable Island

Day number: 009



Date: 10-01-2005


Day number: 010



Date: 11-01-2005


# Sable Island

Day number: 011



Date: 12-01-2005


Day number: 012



Date: 13-01-2005


# Sable Island

Day number: 013



Date: 14-01-2005


Day number: 014



Date: 15-01-2005


# Sable Island

Day number: 015



Date: 16-01-2005


Day number: 016



Date: 17-01-2005


# Sable Island

Day number: 017



Date: 18-01-2005


Day number: 018



Date: 19-01-2005


# Sable Island

Day number: 019



Date: 20-01-2005


Day number: 020



Date: 21-01-2005


# Sable Island

Day number: 021



Date: 22-01-2005


Day number: 022



Date: 23-01-2005


# Sable Island

Day number: 023



Date: 24-01-2005

Day number: 024






Date: 25-01-2005


# Sable Island

Day number: 025



Date: 26-01-2005


Day number: 026



Date: 27-01-2005


# Sable Island

Day number: 027



Date: 28-01-2005


Day number: 028



Date: 29-01-2005


# Sable Island

Day number: 029



Date: 30-01-2005


Day number: 030




Date: 31-01-2005

# Sable Island

Day number: 031




# Sable Island Observatory: Declination (degrees)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Sable Island Observatory: Horizontal Intensity (nT)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Sable Island Observatory: Vertical Intensity (nT)



Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Sable Island Observatory 2005





### Monthly Mean Values for Sable Island Observatory 2005

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	-19° 19.3′	20290 nT	66° 52.9′	19148 nT	-6713 nT	47528 nT	51678 nT

#### Note

i. The values shown here are provisional.